

1-1-1986

Some Probabilities Associated with the Ordering of Unknown Multinomial Cell Probabilities

S. Y. Dennis

Southern Methodist University

Follow this and additional works at: https://scholar.smu.edu/business_workingpapers

 Part of the [Business Commons](#)

This document is brought to you for free and open access by the Cox School of Business at SMU Scholar. It has been accepted for inclusion in Historical Working Papers by an authorized administrator of SMU Scholar. For more information, please visit <http://digitalrepository.smu.edu>.

SOME PROBABILITIES ASSOCIATED WITH THE ORDERING
OF UNKNOWN MULTINOMIAL CELL PROBABILITIES

Working Paper 86-091*

by

S. Y. Dennis

Assistant Professor
of Management Information Sciences
Edwin L. Cox School of Business
Southern Methodist University
Dallas, Texas 75275

*This paper represents a draft of work in progress by the author and is being sent to you for your information and review. Responsibility for the contents rests solely with the author, and it may not be reproduced or distributed without written consent. Please address all correspondence to S. Y. Dennis.

SOME PROBABILITIES ASSOCIATED WITH THE ORDERING
OF UNKNOWN MULTINOMIAL CELL PROBABILITIES

by S. Y. Dennis

Department of Management Information Sciences
Southern Methodist University, Dallas, Texas, USA

SUMMARY

This paper deals with certain probabilities associated with the ordering of the components of a vector of unknown multinomial cell probabilities. The analysis is based upon the assumption that the random multinomial parametric vector is distributed as a Dirichlet distribution. Explicit expressions for the a priori probability that a given event is the most probable or the least probable multinomial event are developed. In addition, recursion formulas are developed that permit the determination of the a priori probability associated with an arbitrary ordering of the unknown cell probabilities.

Some key words: Dirichlet distribution, inverted-Dirichlet distribution, least probable multinomial event, most probable multinomial event, most probable ordering of multinomial events.

1. INTRODUCTION

1.1 Purpose of Research

Many problems concerning the Bayesian analysis of data generating processes involve the multinomial distribution. Unfortunately, sparsity of data may make it impossible to design an optimal sampling procedure for selecting the most or the least probable multinomial event; nevertheless, there is a need to provide some useful information regarding these events for the purposes of supporting decision making. One obvious and useful piece of information is the probability that a given event is the most (least) probable event prior or posterior to sample information; however, explicit expressions for these probabilities are not to be found in the literature. This paper then deals with the development of explicit expressions for these probabilities. In addition, we develop a set of recursion formulas useful for constructing an explicit expression for the probability associated with an arbitrary ordering of the unknown multinomial cell probabilities; recursion formulas for this probability are presented in lieu of an explicit expression in order to minimize notational complexity.

1.2 Background and Preliminaries

Suppose that $Y = (Y_1, \dots, Y_{K+1})$ is a vector of observations having a multinomial distribution with parameter N and random parametric vector $P = (P_1, \dots, P_{K+1})$, where $P_1 + \dots + P_K \leq 1$; hence, the kernel of the likelihood function of the vector of observations is of the form

$$\prod_{j=1}^K p_j^{y_j} (1 - \sum_{j=1}^K p_j)^{N-y_0}$$

where $y_0 = y_1 + \dots + y_K$. By definition a conjugate prior density is a normalized likelihood function; hence, a natural conjugate family of distributions for observations having a multinomial distribution is the Dirichlet family of distributions. Thus, the kernel of the prior density of a Dirichlet distribution with parametric vector $\alpha = (\alpha_1, \dots, \alpha_{K+1})$ is of the form

$$\prod_{j=1}^K p_j^{\alpha_j - 1} (1 - \sum_{j=1}^K p_j)^{\alpha_{K+1} - 1}$$

for any point in the simplex $S_K = \{(p_1, \dots, p_K) : p_j > 0 \text{ for } j = 1, \dots, K \text{ and } p_1 + \dots + p_K \leq 1\}$ in R^K , and zero elsewhere, where the α_j are positive real quantities.

Unfortunately, the evaluation of probability integrals involving the Dirichlet distribution are in general very difficult to carry out directly; primarily because of the simplex constraint placed upon the components of the vector random variable $P = (P_1, \dots, P_{K+1})$. This difficulty, however, is easily overcome through a transformation of random variables suggested by Tiao and Guttman (1965); namely,

$$T: Q_i = P_i (1 - \sum_{j=1}^K P_j)^{-1}, \quad i=1, \dots, K$$

The inverse transformation of T is

$$T^{-1}: P_i = Q_i (1 + \sum_{j=1}^K Q_j)^{-1}, \quad i=1, \dots, K$$

and the Jacobian J of the transformation is found to be

$$J = (1 + \sum_{j=1}^K Q_j)^{-(K+1)}$$

Upon application of the transformation T it follows that the vector random variable $Q = (Q_1, \dots, Q_K)$ is distributed as an inverted-Dirichlet distribution with density function

$$\frac{1}{B(\alpha_1, \dots, \alpha_{K+1})} \prod_{j=1}^K q_j^{\alpha_j - 1} (1 + \sum_{j=1}^K q_j)^{-\sum_{j=1}^{K+1} \alpha_j}$$

for $0 < q_j < \infty$, $j = 1, \dots, K$, in R^K , and zero elsewhere, and where

$$B(\alpha_1, \dots, \alpha_{K+1}) = \Gamma(\alpha_1) \dots \Gamma(\alpha_{K+1}) / \Gamma(\alpha_1 + \dots + \alpha_{K+1}).$$

If we now denote $P_{[1]} \leq \dots \leq P_{[K+1]}$ as the ordered set of unknown cell probabilities associated with the random parametric vector $P = (P_1, \dots, P_{K+1})$, then we can define the most (least) probable multinomial event as the event with the largest (smallest) cell probability.

Utilizing the transformational relationship between the Dirichlet distribution and its multivariate analog, the inverted-Dirichlet distribution, we can write the probability that $P_{[K+1]} = P_k$ as

$$\begin{aligned} & \text{pr}[(1 - \sum_{\substack{j=1 \\ j \neq k}}^{K+1} P_j) \leq P_k] \prod_{\substack{j=1 \\ j \neq k}}^K (P_j \leq P_k) \\ &= \frac{1}{B(\alpha_1, \dots, \alpha_{K+1})} \int_0^\infty \int_0^{q_k} \dots \int_0^{q_k} w_k^{\alpha_k - 1} (1 + \sum_{\substack{j=1 \\ j \neq k}}^K w_j)^{-\sum_{j=1}^{K+1} \alpha_j} \prod_{\substack{j=1 \\ j \neq k}}^K w_j^{\alpha_j - 1} dw_j dw_k \quad (1) \end{aligned}$$

Similarly, we can write the probability that $P_{[1]} = P_k$ as

$$\begin{aligned}
& \text{pr}[(P_k \leq 1 - \sum_{\substack{j=1 \\ j \neq k}}^{K+1} P_j) \cap \bigcap_{\substack{j=1 \\ j \neq k}}^K (P_k \leq P_j)] \\
&= \frac{1}{B(\alpha_1, \dots, \alpha_{K+1})} \int_0^1 \int_{q_k}^\infty \dots \int_{q_k}^\infty w_k^{\alpha_k-1} (1 + \sum_{j=1}^K w_j)^{-\sum_{j=1}^{K+1} \alpha_j} \prod_{\substack{j=1 \\ j \neq k}}^K w_j^{\alpha_j-1} dw_j dw_k \quad (2)
\end{aligned}$$

In like fashion, the probability associated with an arbitrary ordering of the unknown cell probabilities $P = (P_1, \dots, P_{K+1})$, i.e., $P_{(1)} \leq \dots \leq P_{(K+1)}$, can be written as

$$\begin{aligned}
& \text{pr}[\bigcap_{j=1}^K (P_{(j)} \leq P_{(j+1)})] \\
&= \frac{1}{B(\alpha_{(1)}, \dots, \alpha_{(K+1)})} \int_0^1 \int_0^{q_{(K)}} \int_0^{q_{(K-1)}} \dots \int_0^{q_{(2)}} (1 + \sum_{j=1}^K w_{(j)})^{-\sum_{j=1}^{K+1} \alpha_{(j)}} \prod_{j=1}^K w_{(j)}^{\alpha_{(j)}-1} dw_{(j)} \quad (3)
\end{aligned}$$

where the subscripts $((1), \dots, (K+1))$ correspond to exactly one of the $(K+1)!$ permutations of the integers $(1, \dots, K+1)$.

2. THE MOST PROBABLE EVENT

The results of this section are summarized in the following lemma:

Lemma 1. If the vector random variable $P = (P_1, \dots, P_{K+1})$ has a Dirichlet distribution with parametric vector $\alpha = (\alpha_1, \dots, \alpha_{K+1})$, where the α_j are positive integers, then the probability that the k th variate is the largest Dirichlet variate is

$$\begin{aligned} \text{pr}[P_k = \max\{P_1, \dots, P_{K+1}\}] &= 1 - \sum_{i_1} \pi_{i1} + \sum_{i_1 < i_2} \pi_{i1i2} \\ &+ \dots + (-1)^m \sum_{i_1 < \dots < i_m} \pi_{i1 \dots i_m} + \dots + (-1)^K \pi_{1 \dots (k-1)(k+1) \dots K+1} \end{aligned}$$

where $\sum_{i_1 < \dots < i_m}$ denotes the summation over all integers i_1, \dots, i_m , where (i) $1 \leq i_j \leq K+1$, $j = 1, \dots, m$ ($i_j \neq k$) and (ii) $i_1 < \dots < i_m$, and where

$$\pi_{i1 \dots im} = \sum_{x_{i1}=0}^{\alpha_{i1}-1} \dots \sum_{x_{im}=0}^{\alpha_{im}-1} \frac{\Gamma(\alpha_k + \sum_{j=1}^m x_{ij})}{\Gamma(\alpha_k) \prod_{j=1}^m x_{ij}!} \left(\frac{1}{m+1}\right)^{\alpha_k} \prod_{j=1}^m \left(\frac{1}{m+1}\right)^{x_{ij}}$$

To illustrate the application of the above lemma, consider the Dirichlet distributed vector variate $P = (P_1, P_2, P_3, P_4 = 1 - (P_1 + P_2 + P_3))$ having the parametric vector $\alpha = (\alpha_1, \alpha_2, \alpha_3, \alpha_4)$. If the parameters α_j are positive integers, then the probability that the k th variate, where $k=3$, is the largest Dirichlet variate is given by

$$\begin{aligned} \text{pr}[P_3 = \max\{P_1, P_2, P_3, P_4\}] &= 1 - (\pi_1 + \pi_2 + \pi_4) + (\pi_{12} + \pi_{14} + \pi_{24}) - \pi_{124} \\ &= 1 - \sum_{x_1=0}^{\alpha_1-1} \frac{\Gamma(\alpha_3 + x_1)}{\Gamma(\alpha_3) x_1!} \left(\frac{1}{2}\right)^{\alpha_3 + x_1} - \sum_{x_2=0}^{\alpha_2-1} \frac{\Gamma(\alpha_3 + x_2)}{\Gamma(\alpha_3) x_2!} \left(\frac{1}{2}\right)^{\alpha_3 + x_2} \end{aligned}$$

$$\begin{aligned}
& - \sum_{x_4=0}^{\alpha_4-1} \frac{\Gamma(\alpha_3+x_4)}{\Gamma(\alpha_3)x_4!} \left(\frac{1}{2}\right)^{\alpha_3+x_4} + \sum_{x_1=0}^{\alpha_1-1} \sum_{x_2=0}^{\alpha_2-1} \frac{\Gamma(\alpha_3+x_1+x_2)}{\Gamma(\alpha_3)x_1!x_2!} \left(\frac{1}{3}\right)^{\alpha_3+x_1+x_2} \\
& + \sum_{x_1=0}^{\alpha_1-1} \sum_{x_4=0}^{\alpha_4-1} \frac{\Gamma(\alpha_3+x_1+x_4)}{\Gamma(\alpha_3)x_1!x_4!} \left(\frac{1}{3}\right)^{\alpha_3+x_1+x_4} + \sum_{x_2=0}^{\alpha_2-1} \sum_{x_4=0}^{\alpha_4-1} \frac{\Gamma(\alpha_3+x_2+x_4)}{\Gamma(\alpha_3)x_2!x_4!} \left(\frac{1}{3}\right)^{\alpha_3+x_2+x_4} \\
& - \sum_{x_1=0}^{\alpha_1-1} \sum_{x_2=0}^{\alpha_2-1} \sum_{x_4=0}^{\alpha_4-1} \frac{\Gamma(\alpha_3+x_1+x_2+x_4)}{\Gamma(\alpha_3)x_1!x_2!x_4!} \left(\frac{1}{4}\right)^{\alpha_3+x_1+x_2+x_4}
\end{aligned}$$

We will now formally prove Lemma 1, without loss of generality, for the case of $k=K$. The evaluation of probability (1) follows directly from successive application of the following formula established by Taio and Guttman (1965):

$$\begin{aligned}
& \int_0^a (1+x+t)^{-(\alpha+n)} t^{n-1} dt = \\
& (1+x)^{-\alpha} B(\alpha, n) - \sum_{j=0}^{n-1} \binom{n-1}{j} a^j (1+x+t)^{-(\alpha+j)} B(\alpha+j, n-j) \quad (4)
\end{aligned}$$

where n is a positive integer, and $\langle x, \alpha, a \rangle$ are positive quantities, and where $B(u, v) = \Gamma(u)\Gamma(v)/\Gamma(u+v)$.

For simplicity of notation define

$$J_{r \dots K} = \frac{1}{B(\alpha_r, \dots, \alpha_{K+1})} \int_1^\infty \int_0^{q_K} \dots \int_0^{q_K} \left(1 + \sum_{j=1}^K w_j\right)^{-\sum_{j=r}^{K+1} \alpha_j} \prod_{j=r}^K w_j^{\alpha_j-1} dw_j$$

and

$$J_{r \dots K}^{i_1 \dots i_K} = \sum_{x_{i1}=0}^{\alpha_{i1}-1} \dots \sum_{x_{iK}=0}^{\alpha_{iK}-1} \frac{\Gamma(\alpha_{K,k})}{\Gamma(\alpha_K) \prod_{j=1}^k x_{ij}!} \quad (\text{continued on next page})$$

$$\frac{1}{B(\alpha_r, \dots, \alpha_{K,r}, \alpha_{K+1})} \int_1^\infty \int_0^{q_K} \dots \int_0^{q_K} \frac{w_K^{\alpha_{K,k}-1} \prod_{j=r}^{K-1} w_j^{\alpha_j-1} dw_j dw_K}{\sum_{j=r}^{K-1} \alpha_j + \alpha_{K,k} + \alpha_{K+1} \{1 + \sum_{j=r}^{K-1} w_j + (k+1)w_K\}^{j=r}} \quad (5)$$

where i_1, \dots, i_k is an arbitrary sequence of positive integers of length k , $1 \leq k \leq r-1$, such that $1 \leq i_1 < \dots < i_k \leq r-1$, and where $\alpha_{K,k} = \alpha_K + x_{i1} + \dots + x_{ik}$.

For $\alpha_1, \dots, \alpha_K$ positive integers, successive application of integration formula (4) in probability integral (1) yields the following result:

$$J_{1\dots K} = J_K - \sum_{i_1}^{i_1} J_K^{i_1} + \sum_{i_1 < i_2}^{i_1 i_2} J_K^{i_1 i_2} + \dots + (-1)^m \sum_{i_1 < \dots < i_m}^{i_1 \dots i_m} J_K^{i_1 \dots i_m} + \dots + (-1)^K J_K^{1\dots K-1} \quad (6)$$

where $\sum_{i_1 < \dots < i_m}^{i_1 \dots i_m}$ denotes the summation over all integers i_1, \dots, i_m where

(i) $1 \leq i_j \leq K+1$ for $j = 1, \dots, K-1, K+1$ and (ii) $i_1 < \dots < i_m$.

From definitional formula (5) we have that

$$J_K^{i_1 \dots i_k} = \sum_{x_{i1}=0}^{\alpha_{i1}-1} \dots \sum_{x_{ik}=0}^{\alpha_{ik}-1} \frac{\Gamma(\alpha_{K,k})}{\Gamma(\alpha_K) \prod_{j=1}^k x_{ij}!} \left[\frac{1}{B(\alpha_{K,k}, \alpha_{K+1})} \int_1^\infty \frac{q_K^{\alpha_{K,k}-1} dq_K}{\{1 + (1+k)q_K\}^{\alpha_{K,k} + \alpha_{K+1}}} \right] \quad (7)$$

It is well known, see for example Olkin and Sobel (1965), that for any real $r > 0$ and positive integer s

$$\frac{1}{B(r, s)} \int_0^1 t^r (1-t)^{s-1} dt = \frac{1}{B(r, s)} \int_{\theta/(1-\theta)}^\infty u^{r-1} (1+u)^{-(r+s)} du = 1 - I_\theta(r, s) \quad (8)$$

where

$$I_{\theta}(r, s) = \theta^r \sum_{x=0}^{s-1} \frac{\Gamma(r+x)}{\Gamma(r)x!} (1-\theta)^x \quad (9)$$

Thus, using equations (8) and (9) we can write equation (7), upon rearranging, as

$$J_K^{i_1 \dots i_k} = \pi_{i_1 \dots i_k} - \pi_{i_1 \dots i_{k(K+1)}} \quad (10)$$

for $1 \leq k \leq K-1$, where

$$\pi_{i_1 \dots i_k} = \sum_{x_{i1}=0}^{\alpha_{i1}-1} \dots \sum_{x_{ik}=0}^{\alpha_{ik}-1} \frac{\Gamma(\alpha_{K,k})}{\Gamma(\alpha_K) \prod_{j=1}^k x_{ij}!} \left(\frac{1}{k+1}\right)^{\alpha_{K,k}}$$

and where

$$\pi_{i_1 \dots i_{k(K+1)}} = \sum_{x_{i1}=0}^{\alpha_{i1}-1} \dots \sum_{x_{ik}=0}^{\alpha_{ik}-1} \sum_{x_{K+1}=0}^{\alpha_{K+1}-1} \frac{\Gamma(\alpha_{K,k+x_{K+1}})}{\Gamma(\alpha_K) \prod_{j=1}^k x_{ij}! x_{K+1}!} \left(\frac{1}{k+2}\right)^{\alpha_{K,k+x_{K+1}}}$$

Substituting equation (10) into equation (6) yields, upon rearranging, the desired result, i.e.,

$$\begin{aligned} J_{1 \dots K} &= 1 - \sum_{i_1} \pi_{i_1} + \sum_{i_1 < i_2} \pi_{i_1 i_2} + \dots + (-1)^m \sum_{i_1 < \dots < i_m} \pi_{i_1 \dots i_m} \\ &\quad + \dots + (-1)^K \pi_{1 \dots (K-1)(K+1)} \end{aligned}$$

3. THE LEAST PROBABLE EVENT

The results of this section are summarized in the following lemma:

Lemma 2. If the vector random variable $P = (P_1, \dots, P_{K+1})$ has a Dirichlet distribution with parametric vector $\alpha = (\alpha_1, \dots, \alpha_{K+1})$, where the α_j are positive integers, then the probability that the k th variate is the smallest Dirichlet variate is

$$\text{pr}[P_k = \min\{P_1, \dots, P_{K+1}\}] =$$

$$\sum_{x_1=0}^{\alpha_1-1} \dots \sum_{x_{k-1}=0}^{\alpha_{k-1}-1} \sum_{x_{k+1}=0}^{\alpha_{k+1}-1} \dots \sum_{x_{K+1}=0}^{\alpha_{K+1}-1} \frac{\Gamma(\alpha_k + \sum_{j=1}^{K+1} x_j)}{\Gamma(\alpha_k) \prod_{j=1, j \neq k}^{K+1} x_j!} \left(\frac{1}{K+1}\right)^{\alpha_k} \prod_{j=1, j \neq k}^{K+1} \left(\frac{1}{K+1}\right)^{x_j}$$

To demonstrate the application of the above lemma, consider once again the Dirichlet vector variate $P = (P_1, P_2, P_3, P_4 = 1 - (P_1 + P_2 + P_3))$ having the parametric vector $\alpha = (\alpha_1, \alpha_2, \alpha_3, \alpha_4)$. If the parameters α_j are positive integers, then we can write the probability that the k th event, where $k=3$, is the smallest Dirichlet variate as

$$\text{pr}[P_3 = \min(P_1, P_2, P_3, P_4)] =$$

$$\sum_{x_1=0}^{\alpha_1-1} \sum_{x_2=0}^{\alpha_2-1} \sum_{x_4=0}^{\alpha_4-1} \frac{\Gamma(\alpha_3 + x_1 + x_2 + x_4)}{\Gamma(\alpha_3) x_1! x_2! x_4!} \left(\frac{1}{4}\right)^{\alpha_3 + x_1 + x_2 + x_4}$$

We will now formally prove Lemma 2, without loss of generality for the case of $k=K$. The evaluation of probability (2) follows directly from successive application of the following variation of formula (4):

$$\int_a^\infty (1+x+t)^{-(\alpha+n)} t^{n-1} = \sum_{j=0}^{n-1} \binom{n-1}{j} a^j (1+x+a)^{-(\alpha+j)} B(\alpha+j, n-j) \quad (11)$$

where n is a positive integer, and $\langle x, \alpha, a \rangle$ are positive quantities and where $B(u, v) = \Gamma(u)\Gamma(v)/\Gamma(u+v)$.

For simplicity of notation define

$$L_{k+1 \dots K}^{1 \dots k} = \sum_{x_1=0}^{\alpha_1-1} \dots \sum_{x_k=0}^{\alpha_k-1} \frac{\Gamma(\alpha_{K,k})}{\Gamma(\alpha_K) \prod_{j=1}^k x_j!} \cdot \frac{1}{B(\alpha_{k+1}, \dots, \alpha_{K,k}, \alpha_{K+1})} \int_0^1 \int_{q_K}^{\infty} \dots \int_{q_K}^{\infty} \frac{w_K^{\alpha_{K,k}-1} \prod_{j=k+1}^{K-1} w_j^{\alpha_j-1} dw_j dw_K}{\{1 + \sum_{j=k+1}^{K-1} w_j + (k+1)w_K\}^{\alpha_{K,k} + \alpha_{K+1}}}$$

where $\alpha_{K,k} = \alpha_K + x_1 + \dots + x_k$, for $1 \leq k \leq K-1$.

For $\alpha_1, \dots, \alpha_K$ positive integers, successive application of integration formula (11) in probability integral (2) yields the following result:

$$L_{1 \dots K} = L_K^{1 \dots K-1}$$

where

$$L_K^{1 \dots K-1} = \sum_{x_1=0}^{\alpha_1-1} \dots \sum_{x_{K-1}=0}^{\alpha_{K-1}-1} \frac{\Gamma(\alpha_{K,K-1})}{\Gamma(\alpha_K) \prod_{j=1}^{K-1} x_j!} \left(\frac{1}{K}\right)^{\alpha_{K,K-1}} \left\{ \frac{1}{B(\alpha_{K,K-1}, \alpha_{K+1})} \int_0^1 \frac{z^{\alpha_{K,K-1}-1} dz}{(1+z)^{\alpha_{K,K-1} + \alpha_{K+1}}} \right\} \quad (12)$$

It is well known that for any real $r > 0$ and positive integer s

$$\frac{1}{B(r, s)} \int_0^{\theta} \frac{u^{r-1} du}{(1+u)^{r+s}} = I_{\theta/(1+\theta)}(r, s) = \left(\frac{\theta}{1+\theta}\right)^r \sum_{x=0}^{s-1} \frac{\Gamma(r+x)}{\Gamma(r)x!} \left(\frac{1}{1+\theta}\right)^x \quad (13)$$

Hence, using relationship (13) in equation (12) yields the desired result

$$L_{1\dots K} = \sum_{x_1=0}^{\alpha_1-1} \dots \sum_{x_{K-1}=0}^{\alpha_{K-1}-1} \frac{\Gamma(\alpha_{K,K-1}+x_{K+1})}{\Gamma(\alpha_K) \prod_{j=1}^{K-1} x_j! x_{K+1}!} \left(\frac{1}{K+1}\right)^{\alpha_{K,K-1}+x_{K+1}}$$

4. THE MOST PROBABLE ORDERING OF EVENTS

In this section we derive a set of recursion formulas useful for the development of an explicit expression for the probability associated with an arbitrary ordering of the components of a vector random variable having a Dirichlet distribution; specifically the probability implied by equation (3). The derivation of the recursion formulas underlying the evaluation of probability integral (3) follows directly from repeated integration using relationship (4); we once again assume that the components of the parametric vector α are positive integers. For clarity of presentation we drop the parenthesis in subscripts of probability (3) and proceed without loss of generality to evaluate the probability that $P_1 \leq \dots \leq P_{K+1}$.

For simplicity of notation define

$$M_{s\dots K} = \int_0^1 \int_0^{q_K} \int_0^{q_{K-1}} \dots \int_0^{q_{s+1}} \frac{(1 + \sum_{j=s}^K w_j)^{-\sum_{j=1}^{K+1} \alpha_j} (\prod_{j=s}^K w_j^{\alpha_j-1} dw_j)}{B(\alpha_s, \dots, \alpha_{K+1})}, \quad s=1, \dots, K$$

$$M_{s\dots K}^k = \int_0^1 \int_0^{q_K} \int_0^{q_{K-1}} \dots \int_0^{q_{s+1}} \frac{(1 + \sum_{j=s}^K w_j)^{-\sum_{j=1}^{K+1} \alpha_j + x_{s-1}} w_s^{\alpha_s+x_{s-1}-1} (\prod_{j=s+1}^K w_j^{\alpha_j-1} dw_j) dw_s}{B(\alpha_s+x_{s-1}, \alpha_{s+1}, \dots, \alpha_{K+1})}$$

for $s = 2, \dots, K-1$ and $2 \leq k \leq s$,

$$\pi_{r\dots s} = \prod_{j=r}^s \left\{ \sum_{x_j=0}^{\alpha_j+x_{j-1}-1} \frac{\Gamma(\alpha_{j+1}+x_j)}{\Gamma(\alpha_{j+1})x_j!} \left(\frac{1}{j+1}\right)^{\alpha_{j+1}} \left(\frac{j}{j+1}\right)^{x_j} \right\}, \quad s=1, \dots, K-1, \quad r < s \quad (14)$$

and

$$\pi_s = \sum_{x_s=0}^{\alpha_s-1} \frac{\Gamma(\alpha_{s+1}+x_s)}{\Gamma(\alpha_{s+1})x_s!} \left(\frac{1}{2}\right)^{\alpha_{s+1}+x_s}, \quad s=1, \dots, K-1 \quad (15)$$

For $\alpha_1, \dots, \alpha_{K+1}$ positive integers, successive application of formula (4) in equation (3) yields the following recursion formulas:

$$(i) \quad M_{s \dots K} = M_{s+1 \dots K} - \pi_s M_{s+1 \dots K}^2, \quad s=1, \dots, K-1 \quad (16)$$

$$(ii) \quad \pi_{s-(k+1) \dots s-1} M_{s \dots K}^k = \pi_{s-k+1 \dots s-1} M_{s+1 \dots K} \\ - \pi_{s-k+1 \dots s} M_{s+1 \dots K}^{k+2}, \quad s=2, \dots, K-1 \quad (17) \\ 2 \leq k \leq s$$

where

$$M_K = I_{\frac{1}{2}}(\alpha_K, \alpha_{K+1}) = \sum_{x_{K+1}=0}^{\alpha_{K+1}-1} \frac{\Gamma(\alpha_K+x_{K+1})}{\Gamma(\alpha_K)x_{K+1}!} \left(\frac{1}{2}\right)^{\alpha_K+x_{K+1}} \quad (18)$$

and

$$M_K^k = I_{k/k+1}(\alpha_K+x_{K-1}, \alpha_{K+1}) = \sum_{x_{K+1}=0}^{\alpha_{K+1}-1} \frac{\Gamma(\alpha_K+x_{K-1}+x_{K+1})}{\Gamma(\alpha_K+x_{K-1})x_{K+1}!} \left(\frac{k}{k+1}\right)^{\alpha_K+x_{K-1}} \left(\frac{1}{k+1}\right)^{x_{K+1}} \quad (19)$$

using relationship (9).


To illustrate the constructive application of equations (14) and (17), consider the vector random variable $P = (P_1, P_2, P_3, P_4=1-(P_1+P_2+P_3))$ having a Dirichlet distribution with parametric vector $\alpha = (\alpha_1, \alpha_2, \alpha_3, \alpha_4)$. If the parameters α_j are positive integers we can write, using recursion formulas (16)-(17), the probability that $P_1 \leq P_2 \leq P_3 \leq P_4$ as follows:

$$\text{pr}[P_1 \leq P_2, P_2 \leq P_3, P_3 \leq P_4] = M_3 - \pi_2 M_3^2 - \pi_1 M_3 + \pi_{12} M_3^3 \quad (20)$$

Using relationships (14)-(15) and (18)-(19) in probability (20) we have, upon rearranging, the result

$$\begin{aligned}
 \text{pr}[P_1 < P_2, P_2 < P_3, P_3 < P_4] &= \sum_{x_4=0}^{\alpha_4-1} \frac{\Gamma(\alpha_3+x_4)}{\Gamma(\alpha_3)x_4!} \left(\frac{1}{2}\right)^{\alpha_3+x_4} \\
 &- \sum_{x_2=0}^{\alpha_2-1} \sum_{x_4=0}^{\alpha_4-1} \frac{\Gamma(\alpha_3+x_2+x_4)}{\Gamma(\alpha_3)x_2!x_4!} \left(\frac{1}{3}\right)^{\alpha_3+x_2+x_4} - \sum_{x_1=0}^{\alpha_1-1} \frac{\Gamma(\alpha_2+x_1)}{\Gamma(\alpha_2)x_1!} \left(\frac{1}{2}\right)^{\alpha_2+x_1} \sum_{x_4=0}^{\alpha_4-1} \frac{\Gamma(\alpha_3+x_4)}{\Gamma(\alpha_3)x_4!} \left(\frac{1}{2}\right)^{\alpha_3+x_4} \\
 &+ \sum_{x_1=0}^{\alpha_1-1} \frac{\Gamma(\alpha_2+x_1)}{\Gamma(\alpha_2)x_1!} \left(\frac{1}{2}\right)^{\alpha_2+x_1} \sum_{x_2=0}^{\alpha_2+x_1-1} \frac{\Gamma(\alpha_3+x_2)}{\Gamma(\alpha_3)x_2!} \left(\frac{1}{3}\right)^{\alpha_3} \left(\frac{2}{3}\right)^{x_2} \sum_{x_4=0}^{\alpha_4-1} \frac{\Gamma(\alpha_3+x_2+x_4)}{\Gamma(\alpha_3+x_2)x_4!} \left(\frac{3}{4}\right)^{\alpha_3+x_2} \left(\frac{1}{4}\right)^{x_4}
 \end{aligned} \tag{21}$$

The application of the recursion formulas in the above calculation of the probability M_{123} can be visualized with the help of the following tree diagram:



where in equation (20), the root of the tree is set equal to the sum of its terminal branches. The tree is easily extended to yield explicit expressions for any of the probabilities defined by equation (3). For instance, if the root of the tree is M_{1234} then it is easily verified using equations (16) and (17) that the probability that $P_1 < P_2 < P_3 < P_4 < P_5$, where $P_5 = 1 - (P_1 + P_2 + P_3 + P_4)$, is given by

$$\begin{aligned} \text{pr}[P_1 < P_2, P_2 < P_3, P_3 < P_4, P_4 < P_5] &= M_4^{-\pi} 3^{M_4^2} 2^{-\pi} M_4^{\pi} 23^{M_4^3} \\ &- \pi_1^{M_4} 1^{\pi} 3^{M_4^2} 12^{M_4} 123^{M_4^4} \end{aligned} \quad (22)$$

An explicit expression for probability (22) can then be easily obtained using relationships (14)-(15) and (18)-(19) in equation (22).

5. CONCLUSION

We have presented several useful expressions for calculating certain probabilities associated with the ordering of the components of a vector of unknown multinomial cell probabilities. These expressions provide important information in support of the selection of the most (least) probable event or the most probable ordering of multinomial events within a decision framework characterized by a high degree of subjective information and sparsity of data. Admittedly, problems of this ilk occur all too frequently in decision making situations; to this end it is hoped that this study will be of some practical value.

The author thanks Ronald Randles and William Schucany for inspiration and helpful suggestions, respectively, and Lynette Faulkner for her valued and skilled assistance in preparing the manuscript.

REFERENCES

- OLKIN, I. & SOBEL, M. (1965). Integral expressions for tail probabilities of the multinomial and the negative multinomial distributions. *Biometrika* 52, 167-179.
- TIAO, G. G., & GUTTMAN, I. (1965). The inverted-Dirichlet distribution with applications. *J. Am. Statist. Assoc.* 60, 793-805.
(Correction *J. Am. Statist. Assoc.* 60, 1251-1252.)

NOTE: The following is a partial list of papers that are currently available in the Edwin L. Cox School of Business Working Paper Series. A complete list is available upon request from:

Kenneth R. Ferris
Director of Research and Development
Edwin L. Cox School of Business
Southern Methodist University
Dallas, Texas 75275

- 85-908 "Developing Model Strings for Model Managers," by Gary Klein
- 85-909 "Strategic Planning and the Relationship Between Growth, Profitability and Firm Value: An Empirical Study," by Nikhil Varaiya, Roger Kerin, and David Weeks
- 85-110 "Assessment of Multi-Attributed Measurable Value and Utility Functions Via Mathematical Programming," by Gary Klein, Herbert Moskowitz, Sathiadev Mahesh, and A. Ravindran
- 85-111 "Computer Aided Process Structuring Via Mixed Integer Programming," by Gary Klein, P. Beck, and Benn Konsynski
- 85-112 "The Board of Directors, Top Management Compensation, and Shareholder Returns," by J. Kerr, and R.A. Bettis
- 85-113 "An Evaluation of Estimation Procedures for New Product Diffusion Models," by V. Mahajan, C. Mason, V. Srinivasan
- 85-114 "The Impact of Variable Levies: A Discussion of the Costs and Benefits of the European Community's Common Agricultural Policy" by T. Smith
- 85-115 "The Effect of the Bankruptcy Reform Act of 1978 on Small Business loan Pricing," by J. Scott and T. Smith
- 85-116 "The Strategy-Structure Linkage: Conceptual, Methodological and Measurement Issues," by W. Bigler
- 85-117 "A Review of the Framework Oriented Literature on Culture and A Proposed Integration," by W. Bigler
- 85-118 "The Strategy-Culture Linkage: A Procedural Guide," by W. Bigler
- 85-119 "Toward A Theory of Business Level Strategy Research," by W. Bigler
- 85-120 "The Influence of Career Stages on Salespeople's Job Attitudes, Work Perceptions and Performance" by W. Cron and J.W. Slocum
- 85-121 "Stochastic Duration and Dynamic Measure of Risk in Financial Futures," by A. Chen, H. Park, K. Wei
- 86-011 "The Returns to Bidding Firms and the Gains from Corporate Takeovers: A Reexamination," by Nikhil P. Varaiya
- 86-012 "Culture, Key Events and Corporate Social Responsibility," by Lynn A. Isabella
- 86-013 "The Effect of Career Stage and Shared Values on the Interpretation of Key Events," by Lynn A. Isabella
- 86-014 "Systems Analysts Job Responsibilities in the "Profit" vs. the "Not for Profit" Sectors," by Albert Kagan, Marion G. Sobol, and Kevin Quarnstrom

- 86-015 "The Penn Square Bank Failure: Effect on Commercial Bank Security Returns," by John W. Peavy, III, and George H. Hempel
- 86-041 "Career Plateauing: Who's Likely to Plateau?" by John W. Slocum, Jr., William L. Cron, and Linda Yows
- 86-042 "Tax Reform and Housing: Likely Impacts of the Administration Proposal and the House Bill," by David C. Ling and Patric H. Hendershott
- 86-043 "Service Pricing in a Medical Library: A Constrained Goal Programming Approach," by John J. Rousseau
- 86-044 "Fee Assessment in a Medical Library: An Application of the Club Principle," by John J. Rousseau
- 86-051 "Extremal Principle Solutions of Games in Characteristic Function Form: Core, Chebychev and Shapley Value Generalizations," by J. Rousseau and A. Charnes, B. Golany, and M. Keane
- 86-052 "Expert Systems in University Admissions Decisions," by Michael van Breda
- 86-053 "Assessing Trade Show Functions and Performance: An Exploratory Study," by Roger A. Kerin and William L. Cron
- 86-071 "Application of the Means-End Theoretic for Understanding the Cognitive Bases of Performance Appraisal," by James P. Jolly, Thomas J. Reynolds, and John W. Slocum, Jr.
- 86-072 "Understanding the Pricing of Initial Public Offerings," by Andrew J. Chalk and John W. Peavy, III
- 86-073 "Managing Corporate Cultures Through Reward Systems," by Jeffrey Kerr and John W. Slocum, Jr.
- 86-074 "A Causal Analysis of the Impact of Job Performance on the Voluntary Turnover Process," by Ellen F. Jackofsky and John W. Slocum, Jr.
- 86-075 "Career Transitions of superiors and Subordinates," by Suzanne K. Stout, John W. Slocum, Jr., and William L. Cron
- 86-081 "Selecting Joint Venture Partners Is Easy...Almost," by J. Michael Geringer
- 86-082 "Determining Restrictive Goals in Linear Goal Programs," by Philip O. Beck and Gary Klein
- 86-083 "Bicriterion Decision Making Under Uncertainty: An Interactive Approach," by G. Klein, H. Moskowitz, and A. Ravindran

- 86-084 "A Simple Algebraic Estimation Procedure for Innovation
Diffusion Models of New Product Acceptance," by Vijay Mahajan
and Subhash Sharma
- 86-091 "Some Probabilities Associated with the Ordering of Unknown
Multinomial Cell Probabilities," by S. Y. Dennis